

Foods and Arthritis

PhysiciansCommittee
for Responsible Medicine

Millions of people suffer from painful and swollen joints caused by arthritis. Unfortunately, health care providers often don't discuss diet change with patients who have arthritis. This is likely because older research, which tested diets with dairy products, oil, poultry, or meat, showed little benefit.^{1,2} Current research shows that foods can play a substantial role in arthritis.

Different Types of Arthritis

Osteoarthritis is a gradual loss of cartilage and overgrowth of bone in the joints, especially the knees, hips, spine, and fingers. More than 32 million Americans suffer from osteoarthritis, and risk increases with age.³ The condition usually develops gradually and can cause pain and stiffness.

Rheumatoid arthritis (RA), which affects more than 1.3 million Americans, is a more aggressive form of arthritis.⁴ It is an autoimmune disease that causes painful, inflamed joints and can result in permanent damage. RA is one of medicine's mysteries. The disease does not appear in medical reports until the early 1800s, and some suspect that a virus or bacterium may play a role by setting off an autoimmune reaction. Certain genes can also make people more likely to develop RA.

The Role of Diet

Diet and RA. While not all research has found a connection, multiple studies show that dietary changes can help relieve RA symptoms.⁵ For example, one study looked at the effects of a very-low-fat vegan diet on people with RA. After only four weeks,



people had less morning stiffness, RA pain, joint tenderness, and joint swelling.⁶ A study published in *Rheumatology* found a gluten-free, vegan diet improved the signs and symptoms of RA.⁷ And a systematic review concluded that fasting followed by a vegetarian or vegan diet might be useful in the treatment of RA.⁸

Weight loss can also help. Studies have shown that people with RA who are overweight tend to have worse outcomes than those who are at a healthy weight.^{9,10} In fact, one study found that

While not all research has found a connection, **multiple studies** show that **dietary changes** can help relieve **RA symptoms**.⁵

overweight people with RA who lost 11 or more pounds were three times more likely to have improvement in their disease than those who lost less than 11 pounds.¹¹

Finally, research suggests that certain foods can trigger RA symptoms in some people.¹² Animal products like meat, eggs, and dairy appear to be common triggers. Some healthy foods can be triggers too, however, such as citrus, tomatoes, and corn, to name a few.¹³ Not everyone will have food triggers, though, and when present, the exact foods that trigger symptoms are often unique to that person.¹³ That is, one person's trigger foods could be eaten by someone else with no problems.

Diet and osteoarthritis. People with osteoarthritis can also benefit from dietary changes. Research on more than 6,000 people revealed that the more fiber people ate, the less likely they were to have osteoarthritis symptoms.^{14,15} For overweight people, weight loss can also help with osteoarthritis. Finally, a 2015 study found that people eating a whole food, plant-based diet significantly decreased their osteoarthritis pain—in just two weeks. By the end of the six-week study, they reported more energy and better physical functioning, too.¹⁶

How Might Plant-Based Diets Help?

Vegan diets are often lower in fat and have a healthier fat profile than diets containing animal products. For example, fruits, vegetables, whole grains, and beans typically have a healthful ratio of omega-6 and omega-3 fats. Vegan diets are also generally low in saturated fat. These changes in the amount and types of fat are anti-inflammatory. Weight loss that results from eating a plant-based diet also fights inflammation.

Another potential problem with diets containing animal products is that meat can supply an overload of iron. Too much iron triggers production of dangerous free radicals—basically causing “rusting” inside the body. Excess free radicals can damage joints and contribute to heart disease and cancer.¹⁷

Vegetables and beans, on the other hand, contain ample iron, but in a form the body absorbs only when needed. Moreover, fruits and vegetables like citrus fruits, peppers, and leafy greens contain antioxidants like vitamins C and E, to keep free radicals in check.

Food Triggers for Arthritis

Specific foods may contribute to joint pain for some, particularly for those with RA.¹²

If you’d like to find out if any foods trigger your arthritis pain, here’s how to get started. For two weeks:

- Enjoy an abundance of foods from the pain-safe list.
- Avoid the common triggers completely.
- Foods that are not on either list can be eaten freely.

If your diet change makes your pain disappear or improve, the next step is to find out which foods are your triggers. To do this, simply reintroduce the eliminated foods one at a time, every two days, to see whether any symptoms result. If a food triggers symptoms for you, wait until your symptoms improve before trying the next food. Animal products are best left permanently off your plate.

Pain-Safe Foods

Pain-safe foods virtually never contribute to arthritis, headaches, or other painful conditions. These include:

- Rice
- Cooked green vegetables
- Cooked orange vegetables
- Cooked yellow vegetables
- Cooked or dried noncitrus fruits

Possible Pain Triggers

Common food triggers can cause pain in susceptible people. Certain drinks and additives are also common triggers, including alcoholic beverages (especially red wine), caffeinated drinks (coffee, tea, and colas), monosodium glutamate (MSG), aspartame (NutraSweet), and nitrites (processed meat such as hot dogs and bacon).

Here are some common food triggers.¹⁸⁻²⁰

- Meat, dairy, and eggs
- Wheat, barley, and rye
- Corn
- Nuts and peanuts
- Soy
- Citrus fruits
- Chocolate
- Nightshade vegetables: tomatoes, white potatoes, peppers (except black pepper), and eggplant
- Oats
- Sugar (cane)
- Apples
- Onions
- Celery
- Chickpeas
- Bananas
- Sweet potatoes
- Nutritional yeast

One important note: If you suspect wheat or gluten may be triggering your symptoms, please check with your health care provider about whether you should be screened for celiac disease before eliminating gluten-containing foods from your diet.

Other Approaches

The first line of defense in arthritis is a lower-fat, plant-based diet with plenty of fruits and vegetables, a vitamin B12 supplement, and medications as prescribed by your health care provider.

However, some studies suggest that supplementing with the omega-3 fats DHA and EPA may help people with RA. If you wish to try omega-3s, look for a vegan DHA/EPA supplement (made from algae) with no more than 2 grams of DHA/EPA combined. High omega-3 intakes can increase bleeding risk and interact with some medications, so check with your health care provider before using.²¹

Small studies also suggest that curcumin, found in turmeric, may help ease osteoarthritis pain. Until more research is available, season food such as chili, soup, and curries with turmeric instead of using supplements.²² Be sure to add a pinch of black pepper if you like it. It can help you absorb more curcumin from the turmeric.

This fact sheet is not intended as a comprehensive program for arthritis. If you have arthritis, please consult with your health care provider about which interventions are appropriate for you.

References

1. Panush RS, Carter RL, Katz P, Kowsari B, Longley S, Finnie S. Diet therapy for rheumatoid arthritis. *Arthritis Rheum*. 1983;26(4):462-471. doi: 10.1002/art.1780260403
2. Lithell H, Bruce A, Gustafsson IB, et al. A fasting and vegetarian diet treatment trial on chronic inflammatory disorders. *Acta Derm Venereol*. 1983;63(5):397-403.
3. Centers for Disease Control and Prevention. Osteoarthritis (OA). Centers for Disease Control and Prevention website. <https://www.cdc.gov/arthritis/basics/osteoarthritis.htm>. Accessed April 30, 2021.
4. Helmick CG, Felson DT, Lawrence RC, et al. Estimates of the prevalence of arthritis and other rheumatic conditions in the United States. Part I. *Arthritis Rheum*. 2008;58(1):15-25. doi: 10.1002/art.23177
5. Smedslund G, Byfuglien MG, Olsen SU, Hagen KB. Effectiveness and safety of dietary interventions for rheumatoid arthritis: a systematic review of randomized controlled trials. *J Am Diet Assoc*. 2010;110(5):727-735. doi: 10.1016/j.jada.2010.02.010
6. McDougall J, Bruce B, Spiller G, Westerdahl J, McDougall M. Effects of a very low-fat, vegan diet in subjects with rheumatoid arthritis. *J Altern Complement Med*. 2002;8(1):71-75. doi: 10.1089/107555302753507195
7. Hafström I, Ringertz B, Spångberg A, et al. A vegan diet free of gluten improves the signs and symptoms of rheumatoid arthritis: the effects on arthritis correlate with a reduction in antibodies to food antigens. *Rheumatology (Oxford)*. 2001;40(10):1175-1179. doi: 10.1093/rheumatology/40.10.1175
8. Müller H, de Toledo FW, Resch KL. Fasting followed by vegetarian diet in patients with rheumatoid arthritis: a systematic review. *Scand J Rheumatol*. 2001;30(1):1-10. doi: 10.1080/030097401750065256
9. Gremese E, Carletto A, Padovan M, et al. Obesity and reduction of the response rate to anti-tumor necrosis factor α in rheumatoid arthritis: an approach to a personalized medicine. *Arthr Care Res*. 2013;65(1):94-100. doi: 10.1002/acr.21768
10. Sandberg ME, Bengtsson C, Källberg H, et al. Overweight decreases the chance of achieving good response and low disease activity in early rheumatoid arthritis. *Ann Rheum Dis Lond*. 2014;73(11):2029-2033. doi: 10.1136/annrheumdis-2013-205094
11. Kreps DJ, Halperin F, Desai SP, et al. Association of weight loss with improved disease activity in patients with rheumatoid arthritis: a retrospective analysis using electronic medical record data. *Int J Clin Rheumatol*. 2018;13(1):1-10. doi: 10.4172/1758-4272.1000154
12. Alwarith J, Kahleova H, Rembert E, et al. Nutrition interventions in rheumatoid arthritis: the potential use of plant-based diets. a review. *Front Nutr*. 2019;6:141-152. doi: 10.3389/fnut.2019.00141
13. Darlington LG, Ramsey NW. Review of dietary therapy for rheumatoid arthritis. *Br J Rheumatol*. 1993;32(6):507-514. doi: 10.1093/rheumatology/32.6.507
14. Dai Z, Lu N, Niu J, Felson DT, Zhang Y. Dietary fiber intake in relation to knee pain trajectory. *Arthritis Care Res (Hoboken)*. 2017;69(9):1331-1339. doi: 10.1002/acr.23158
15. Dai Z, Niu J, Zhang Y, Jacques P, Felson DT. Dietary intake of fibre and risk of knee osteoarthritis in two US prospective cohorts. *Ann Rheum Dis*. 2017;76(8):1411-1419. doi: 10.1136/annrheumdis-2016-210810
16. Clinton CM, O'Brien S, Law J, Renier CM, Wendt MR. Whole-foods, plant-based diet alleviates the symptoms of osteoarthritis. *Arthritis*. 2015;2015:708152-708161. doi: 10.1155/2015/708152
17. Merry P, Grootveld M, Lunec J, Blake DR. Oxidative damage to lipids within the inflamed human joint provides evidence of radical-mediated hypoxic-reperfusion injury. *Am J Clin Nutr*. 1991;53(1 Suppl):362S-369S. doi: 10.1093/ajcn/53.1.362S
18. Lidén M, Kristjánsson G, Valtysdóttir S, Venge P, Hällgren R. Self-reported food intolerance and mucosal reactivity after rectal food protein challenge in patients with rheumatoid arthritis. *Scand J Rheumatol*. 2010;39(4):292-298. doi: 10.3109/03009740903379630
19. Karatay S, Erdem T, Kiziltunc A, et al. General or personal diet: the individualized model for diet challenges in patients with rheumatoid arthritis. *Rheumatol Int*. 2006;26(6):556-560. doi: 10.1007/s00296-005-0018-y
20. Darlington LG, Ramsey NW. Review of dietary therapy for rheumatoid arthritis. *Br J Rheumatol*. 1993;32(6):506-514. doi: 10.1093/rheumatology/32.6.507
21. National Institutes of Health. Omega-3 Fatty Acids. National Institutes of Health Office of Dietary Supplements website. <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/>. Accessed May 10, 2021.
22. National Institutes of Health. Turmeric. National Center for Complementary and Integrative Health website. <https://nccih.nih.gov/health/turmeric/ataglance.htm>. Accessed May 10, 2021.